

*Alissonotum pauper* (BURMEISTER, 1847) クロマルコガネ

*Heteronychus pauper* BURMEISTER, Handb. Ent. V, p. 94, 1847 (Philippines).

Distr. Japan (Okino erabu Is., Tokara Is.), Taiwan, China, Philippines, Indo-China.

Subfamily EUCHIRINAE テナガコガネ亞科

Genus *Cheirotonus* HOPE, 1841 テナガコガネ属

*Cheirotonus* HOPE, 1841, Trans. Linn. Soc. (17), p. 594.

(type-species: *Cheirotonus macleayi* HOPE, 1841)

*Cheirotonus jambar* Y. KUROSAWA, 1984 ヤンバルテナガコガネ

*Cheirotonus jambar* Y. KUROSAWA, Bull. Natn. Sci. Mus. Tokyo, Ser. A. 10(2):73-78, Fif. 1-3, 1984 (N. Okinawa Is.).

Distr. Japan (N. Okinawa Is.).

Subfamily CETONIINAE ハナムグリ亞科

Tribe HETERORRHININI カナブン族

Genus *Cosmiomorpha* SAUNDERS, 1852 チヤイロカナブン属

*Cosmiomorpha* SAUNDERS, 1852, Trans. ent. Soc. London (n. s.) II, p. 28.

(type-species: *Cosmiomorpha modesta* SAUNDERS, 1852)

Subgenus *Microcosmiomorpha* MIKŠIĆ

*Microcosmiomorpha* MIKŠIĆ, 1974, Rev. suis Zool. LXXXI. 4, p. 770.

(type-species: *Cosmiomorpha setulosa* WESTWOOD, 1854).

*Cosmiomorpha* (*Microcosmiomorpha*) *similis* FAIRMAIRE, 1899 チヤイロカナブン

*Cosmiomorpha similis* FAIRMAIRE, Ann. Soc. Ent. Fr. LXVII, p. 621, 1899.

Distr. Japan, Taiwan.

subsp. *nigra* NIJIMA et KINOSHITA, 1927 チヤイロカナブン

*Cosmiomorpha similis* FAIRMAIRE var. *nigra* NIJIMA et KINOSHITA, Res. Bull. Coll. Exp. Forest Coll. Agr. Hokkaido Imp. Univ. Japan IV:69, 1927 (Ishigaki-jima).

Distr. Japan (Ishigaki-jima, Iriomote-jima, Obama-jima).

subsp. *miyakoana* NOMURA, 1964 ミヤコチヤイロカナブン

*Cosmiomorpha similis miyakoana* NOMURA, Tōhō Gakuho (14):70, 1964 (Miyako Is.).

Distr. Japan (Miyako-jima, Irabu-jima).

subsp. *yonakuniana* NOMURA, 1964 ヨナクニチヤイロカナブン

*Cosmiomorpha similis yonakuniana* NOMURA, Ent. Rev. Japan 16(2):57, pl. 8, f. 13, 1864 (Yonaku-ni Is.).

Distr. Japan (Yonaguni-jima).

Genus *Rhomorrhina* HOPE, 1837 アオカナブン属  
*Rhomorrhina* HOPE, 1837, Coleop. Man. I, p. 201.  
 (type-species: *Cetonia resplendens* SWARTZ, 1817)

*Rhomorrhina hamai* NOMURA, 1964 イシガキアオカナブン

*Rhomorrhina hamai* NOMURA, Ent. Rev. Japan 16(2):57-58, pl. 8, f. 15, 1964 (Ishigaki Is.).

Distr. Japan (Ishigaki Is., Iriomote Is.).

*Rhomorrhina polita* WATERHOUSE, 1875 クロカナブン

*Rhomorrhina polita* WATERHOUSE, Trans. ent. Soc. London, p. 113, 1875 (Yokohama).

Distr. Japan (Honshu, Sado, Izu-Niijima, Shikoku, Kyushu, Hekurajima, Tanegashima, Yakushima).

*Rhomorrhina unicolor* MOTSCHULSKY, 1861 アオカナブン

*Rhomorrhina unicolor* MOTSCHULSKY, Etud. Ent., X, p. 8, 1861.

Distr. Japan (Hokkaido, Honshu, Sado Is., Shikoku, Kyushu, Hirato-jima, Goto Is., Tsushima, Tokunoshima), Korea.

Genus *Pseudotorynorrhina* MIKŠIĆ, 1967 カナブン属

*Pseudotorynorrhina* MIKŠIĆ, Entom. Abhand. Statt. Mus. Tierk. Dresden 35(3), p. 309, 1967.  
 (type-species: *Rhomorrhina japonica* HOPE, 1841)

*Pseudotorynorrhina japonica* (HOPE, 1841) カナブン

*Rhomorrhina japonica* HOPE, Trans. ent. Soc. London, 1841, p. 64 (Japan).

*Rhomorrhina ignita* NONFRIED, Deut. Ent. Zeit., p. 90, 1890.

*Rhomorrhina japonica* var. *clypeata* BURMEISTER, Handb. Ent. III, p. 199, 780, 1842.

*Rhomorrhina glauca* THOMSON, Typi Cet., p. 9, 1878.

*Rhomorrhina squaligera* THOMSON, Typi Cet., p. 9, 1878.

Distr. Japan (Honshu, Sado Is., Izu Is.-Nii-jima, Kozu-jima, Oshima, Shikoku, Kyushu, Yakushima), Saisyuto, Korea.

Tribe DIPLOGNATHINI マダラハナムグリ族

Genus *Anthracophora* BURMEISTER, 1842 マダラハナムグリ属

*Anthracophora* BURMEISTER, 1842, Handb. Entom. III, p. 623  
 (type-species: *Anthracophora rusticola* BURMEISTER 1842).

*Anthracophora rusticola* BURMEISTER, 1842 アカマダラハナムグリ

*Anthracophora rusticola* BURMEISTER, Handb. Ent. III, p. 624, 1842.

Distr. Japan (Honshu, Sado, Shikoku, Kyushu, Yakushima).

## Tribe CETONIINI ハナムグリ族

Genus *Glycyphana* BURMEISTER, 1842 クロハナムグリ属

*Glycyphana* BURMEISTER, Handb. Ent. III, p. 346, 1842.  
(type-species: *Cetonia festiva* FABRICIUS, 1792).

*Glycyphana fulvistemma* MOTSCHULSKY, 1860 クロハナムグリ

*Glycyphana fulvistemma* MOTSCHULSKY, in Schrenk's Reise Amurländer, p. 135, 1860.

*Euryomia sieboldi* SNELLEN VAN VOLLENHOVEN, 1864, Tijdschr. Ent., 7, p. 158.

*Glycyphana fulvistemma* ab. *nagoyanus* TAKAGI, 1936, Insect World Gifu 40(467):256.

Distr. Japan (Hokkaido, Honshu, Shikoku, Kyushu, Tsushima), Saisyu-to, Korea, Mongolia, Dauria, Taiwan.

*Glycyphana gracilis* SAWADA, 1942 ホソコハナムグリ

*Glycyphana gracilis* SAWADA, Zool. Mag. 56(6):240-241, 1942.

Distr. Japan, Taiwan, Indochina Pen.

subsp. *viridis* SAWADA, 1942 ホソコハナムグリ

*Glycyphana gracilis viridis* SAWADA, Zool. Mag. 56(6):241-242, 1942 (Nara, Tosa, Kagoshima).

Distr. Japan (Honshu, Shikoku, Kyushu, Yakushima).

Genus *Cetonia* FABRICIUS, 1775 ハナムグリ属

*Cetonia* FABRICIUS, Syst. Ent. I, p. 52, 1775.  
(type-species: *Scarabaeus auratus* LINNÉ)

Subgenus *Eucetonia* SCHOCH, 1894

*Eucetonia* SCHOCH, Mitth. Schweiz. Ent. IX, p. 211, 1894.  
(type-species: *Glycyphana viridiopaca* MOTSCHULSKY, 1860)

*Cetonia (Eucetonia) pilifera* (MOTSCHULSKY, 1860) ハナムグリ

*Glycyphana pilifera* MOTSCHULSKY, Etud. Ent. 1860, p. 15.

Distr. Japan (Hokkaido, Honshu, Izu-Rishima, Okino-shima, Shikoku, Kyushu, Goto Is.), Saisyu-to.

subsp. *izuensis* NOMURA, 1969

*Cetonia (Eucetonia) pilifera izuensis* NOMURA, Entom. Rev. Japan 21(2):75, 1969 (Ōshima, Toshima, Niijima, Shikine-jima, Kozu-jima, Mikura-jima, Izu Is.).

Distr. Japan (Izu Is.).

*Cetonia (Eucetonia) roelofsi* HAROLD, 1880 アオハナムグリ

*Cetonia roelofsi* HAROLD, Bull. Soc. ent. Belg., 1880, p. 5.

*Gametis aygyrostica* BURMEISTER, 1842, Handb. Ent., 3, p. 360.

*Cetonia roelofsi* ab. *nitidula* REITTER, 1898, Verh. naturw. Ver. Brunn, 37, p. 55.

*Cetonia roelofsi* ab. *villifrons* REITTER, 1898, Verh. naturw. Ver., Brunn, 37, p. 55.

*Cetonia bodymeyeri* CURTI, Entom. Mitteil. IV, p. 20, 1915.

*Cetonia roelofsi gotoana* NAKANE, 1964, Akitu, Kyoto, 11(4):48.

- Distr. Japan (Hokkaido, Honshu, Shikoku, Kyushu, Yakushima), S. Kurile Is.  
 subsp. *iijima* K. SAKAI, 1996  
*Cetonia (Eucetonia) roelofsi iijimai* K. SAKAI, Gekkan Mushi, (309):2, 10–11, pl. 1, fig. 1–4, 1996  
 (Nishino-shima, Oki Is.).  
 subsp. *tane* NAKANE, 1993  
*Cetonia roelofsi tane* NAKANE, LAMELLICORNIA (9):2, 1993 (Is. Tanegashima).  
 Distr. Japan (Tanegashima).

Genus *Protaetia* BURMEISTER, 1842 シロテンハナムグリ属  
*Protaetia* BURMEISTER, 1842, Handb. Entom. III, p. 472.  
 (type species: *Cetonia anovittata* CHEVROLAT, 1841)

Subgenus *Liocola* C. G. THOMSON, 1859  
*Liocola*, C. G. THOMSON, 1859, Skand. Col. I, p. 77.  
 (type-species: *Cetonia lugubris* HERBST, 1786).

- Protaetia (Liocola) brevitarsis* (LEWIS, 1879) シラホシハナムグリ  
*Cetonia brevitarsis* LEWIS, 1879, Ann. Mag. Nat. Hist. (5)IV, p. 463.  
 Distr. Japan, Korea, E. Siberia.  
 subsp. *brevitarsis* (LEWIS, 1879) シラホシハナムグリ  
*Cetonia brevitarsis* LEWIS, 1879, Ann. Mag. Nat. Hist. (5)IV, p. 463.  
 Distr. Japan (Hokkaido, Honshu, Sado, Shikoku, Kyushu, Hekura-jima, Yakushima), S. Kurile.  
 subsp. *seulensis* (KOLBE, 1886) チョウセンシラホシハナムグリ  
*Cetonia seulensis* KOLBE, 1886, Arch. Naturgesch LII, p. 194, tab. 11, fig. 29.  
 Distr. Japan (Tsushima), Korea, China, Siberia.

- Protaetia (Liocola) cataphracta* ARROW, 1913 ムラサキツヤハナムグリ  
*Protaetia cataphracta* ARROW, 1913, Ann. Mag. Nat. Hist. (8)XII, p. 405.  
 Distr. Japan (Hokkaido, Honshu, Tobishima, Sado, Izu Is.–Mikurajima, Shikoku, Kyushu, Yakushima).

- Protaetia (Liocola) formosana* (MOSER, 1910) タイワンツヤハナムグリ  
*Liocola formosana* MOSER, 1910, Ann. Soc. Ent. Belg. LIV, p. 357.  
 Distr. Japan, Taiwan.  
 subsp. *nozatoi* Y. MIYAKE, 1986 ヨナグニツヤハナムグリ  
*Protartia formosana nozatoi* Y. MIYAKE, 1986, LAMELLICORNIA (2):5 (Yonaguni Is.).  
 Distr. Japan (Yonaguni Is.).

- Protaetia (Liocola) insperata* (LEWIS, 1879) ミヤマオオハナムグリ  
*Cetonia insperata* LEWIS, 1879, Ann. Mag. Nat. Hist., (5)4, p. 463. (Yeso, Vries Island, Chuzenji, Nikko, Wada-toge).  
 Distr. Japan (Hokkaido, Honshu, Kyushu).

- Protaetia (Liocola) miyakoensis* (NIIJIMA et KINOSHITA, 1923) ミヤコオオハナムグリ  
*Liocola miyakoensis* NIIJIMA et KINOSHITA, Res. Bull. Coll. Exp. Forest. Coll. Agr. Hokkaido

Imp. Univ. II, 2, 1923, pp. 170, 233, Taf. II, f. 6, Taf. V, f. 14, Taf. VII, f. 17 (Miyako Is., Tosa in Shikoku).

Distr. Japan (Miyako-jima, Okinawa Is., Irabu-jima).

Subgenus *Tomentoprotetaetia* Mirešić, 1987

*Tomentoprotetaetia* Mirešić, Cetoniinae der Pal. und Orient. Region (4):444-445, 1987.  
(type-species: *Cetonia (Protaetia) culta* WATERHOUSE, 1879).

*Protaetia (Tomentoprotetaetia) culta* (WATERHOUSE, 1879) カバイロハナムグリ  
*Cetonia (Protaetia) culta* WATERHOUSE, 1879, Ann. Mag. Nat. Hist. (5)IV, p. 247.  
*Potosia grisea* NIIZIMA et KINOSHITA, 1923, Res. Bull. Coll. exp. Hokkaido Imp. Univ., II, p. 234.  
*Protaetia culta yaeyamana* NOMURA, 1965, Entom. Rev. Japan XVIII(1):9.  
Distr. Japan (Ishigaki-jima, Iriomote-jima, Yonakuni-jima), Taiwan.

Subgenus *Potosia* Mulsant & Rey, 1871

*Cetonia* subg. *Potosia* Mulsant & Rey, 1871, Col. France Lamell. ed. 2, p. 669.  
(type-species: *Cetonia cuprea* Fabricius, 1775, Syst. Ent. p. 48.)

*Protaetia (Potosia) exasperata* (FAIRMAIRE, 1898) オオシマアオハナムグリ  
*Cetonia exasperata* FAIRMAIRE, Ann. Soc. ent. France 67:386, 1898 (Oshima).  
Distr. Japan.  
subsp. *akitai* MASUMOTO et K. SAKAI, 1988 クメジマアオハナムグリ  
*Protaetia exasperata akitai* MASUMOTO et K. SAKAI, Gekkan Mushi (208):3-5, pl. 2, fig. 1-10, 1988  
(Kumejima, Okinawa).  
Distr. Japan (Kume-jima, Is. Okinawa).  
subsp. *erabuana* NOMURA, 1964 クチノエラブアオハナムグリ  
*Protaetia exasperata erabuana* NOMURA, 1964, Entom. Rev. Japan 16(2):58, pl. 8, f. 16 (Kuchinoera-  
bu Is.).  
Distr. Japan (Kuchinoerabu Is.).  
subsp. *exasperata* (FAIRMAIRE, 1898) オオシマアオハナムグリ  
*Cetonia exasperata* FAIRMAIRE, Ann. Soc. ent. France 67:386, 1898 (Oshima).  
Distr. Japan (Amami-Oshima, Tokunoshima, Obaba-jima).  
subsp. *iheyana* K. SAKAI, 1989 イヘヤアオハナムグリ  
*Protaetia exasperata iheyana* K. SAKAI, Gekkan Mushi (218):18-19, 1989 (Iheyajima Is.).  
Distr. Japan (Iheya-jima).  
subsp. *nakana* NAKANE, 1956 トカラアオハナムグリ  
*Protaetia (Potosia) exasperata nakana* NAKANE, Sci. Rep. Saikyo Univ. (Nat. Sci. Liv. Sci.).  
2(2):61, 1956, pl. I, figs. 3, 5, II; pl. II, Fig. 7 (Nakanoshima).  
Distr. Japan (Nakanoshima, Kuchinoshima).  
f. *tsuzimotoi* NAKANE, 1956, Sci. Rep. Saikyo Univ. Kyoto (A)2(2):61.  
subsp. *satoi* NAKANE, 1956 オキナワアオハナムグリ  
*Protaetia (Potosia) exasperata satoi* NAKANE, 1956, Sci. Rep. Saikyo Univ. (Nat & Liv. Sci.) 2(2)A:  
61, pl. I, Fig. 1, 4, pl. II, Fig. 8.  
Distr. Japan (Okinawa, Tokashiki, Ishigaki, Iriomote Is.).

subsp. *suwanoseana* NOMURA, 1964 スワノセアオハナムグリ  
*Protaetia exasperata suwanoseana* NOMURA, Tōhō Gakuho (14):69, 1964 (Suwanose-jima-Tokara Is.).

Distr. Japan (Suwanose Is., Akuseki Is.-Tokara Is.).  
 subsp. *uenoi* NOMURA, 1960 エラブアオハナムグリ

*Protaetia exasperata uenoi* NOMURA, 1960, Tōhō Gakuho (10):75 (Okinoerabu-jima).  
 Distr. Japan (Okinoerabu-jima).

*Protaetia (Potosia) famelica* (JANSON, 1879) シナハナムグリ

*Cetonia famelica* JANSON, 1879, Cist. Ent. II, p. 539.

Distr. Japan, Saishu-to, Korea, China, E. Siberia.

subsp. *amurensis* (HEYDEN, 1886) シナハナムグリ

*Cetonia amurensis* HEYDEN, 1886, Deutsche Entom. Zeitschr. XXX, p. 285.  
 Distr. Japan (Tsushima), Saishu-to, Korea, E. Siberia.

#### Subgenus *Heteroprotetaetia* MIKŠIĆ, 1963

*Protaetia* subg. *Heteroprotetaetia* MIKŠIĆ, 1963, Entom. Abhand. Staat. Mus. Tierk. Dresden, XXIX, 4, p. 355.  
 (type-species: *Cetonia fusca* HERBST, 1790)

*Protaetia (Heteroprotetaetia) fusca* (HERBST, 1790) ハイイロハナムグリ

*Cetonia fusca* HERBST, 1790, Natursyst Käf. III, p. 257, tab. 32, fig. 4.

*Protaetia taiwana* NIIZIMA et KINOSHITA, 1923, Res. Bull. Coll. Exp. Forest, Coll. Agr. Hokkaido Imp. Univ., II, 2, pp. 176, 235, tab. II, fig. 4.

Distr. Japan (Ogasawara Is., Kimajima-Miyako Is., Irabu-jima), E. S. Asia, Micronesia.

#### Subgenus *Pseudocalopotosia* MIKŠIĆ, 1966

*Protaetia* subgenus *Pseudocalopotosia* MIKŠIĆ, 1966, Zeitschr. Arbeitsgemein. öster. Entom. XVIII, 1, p. 6.  
 (type-species: *Cetonia ishigakia* FAIRMAIRE, 1898)

*Protaetia (Pseudocalopotosia) ishigakia* (FAIRMAIRE, 1898) イシガキシロテンハナムグリ

*Cetonia ishigakia* FAIRMAIRE, 1898, Bull. Soc. Ent. France, p. 14 (Ishigaki Is.).

Distr. Japan.

subsp. *ishigakia* (FAIRMAIRE, 1898) イシガキシロテンハナムグリ

*Cetonia ishigakia* FAIRMAIRE, Bull. Soc. Ent. France, 1898, p. 14 (Ishigaki Is.).

*Protaetia ishigakia iromoteana* NOMURA, 1964, Entom. Rev. Japan 16(2):59, pl. 8, fig. 19.

Distr. Japan (Ishigaki-jima, Iriomote-jima, Hateruma-jima, Obama-jima).

subsp. *kikaiana* NAKANE, 1960 キカイシロテンハナムグリ

*Protaetia ishigakia kikaiana* NAKANE, 1960, Entom. Rev. Japan 12:6 (Kikaijima).

Distr. Japan (Kikaijima).

subsp. *miyakoana* Y. KUROSAWA, 1959 ミヤコシロテンハナムグリ

*Protaetia ishigakia miyakoana* Y. KUROSAWA, 1959, Bull. Nat. Sci. Mus. Tokyo 44:339, fig. 6 (Miyakojima).

Distr. Japan (Miyako-jima, Irabu-jima).

- subsp. *okinawana* Y. KUROSAWA, 1959 オキナワシロテンハナムグリ  
*Protaetia ishigakia okinawana* Y. KUROSAWA, 1959, Bull. Nat. Sci. Mus. Tokyo 44:339, fig.C. (Okinawa).  
Distr. Japan (Okinawa-honto, Iejima, Yoronto, Okinoerabujima, Irabujima).  
subsp. *yonakuniana* NOMURA, 1964 ヨナクニシロテンハナムグリ  
*Protaetia ishigakia yonakuniana* NOMURA, 1964, Entom. Rev. Japan 16(2):59, pl.8, fig.20 (Yonakuni Is.).  
Distr. Japan (Yonaguni-jima).

Subgenus *Calopotosia* REITTER, 1898  
*Potosia Calopotosia* REITTER, 1898, Best. Tab. XXXVIII, p. 49.  
(type-species: *Cetonia orientalis* GORY-PERCHERON, 1833)

- Protaetia (Calopotosia) lewisi* (JANSON, 1888) リュウキュウオオハナムグリ  
*Cetonia lewisi* JANSON, 1888, Ann. Mag. Nat. Hist. (6) I, p. 194.  
Distr. Japan.  
subsp. *leachi* Y. KUROSAWA, 1859 アマミオオハナムグリ  
*Protaetia lewisi leachi* Y. KUROSAWA, 1959, Bull. Nat. Sci. Mus. Tokyo 44:337, fig.f. (Amami-Oshima).  
Distr. Japan (Amami-Oshima, Tokuno-shima).  
subsp. *lewisi* (JANSON, 1888) リュウキュウオオハナムグリ  
*Cetonia lewisi* JANSON, Ann. Mag. Nat. Hist. (6) I, 1888, p. 194 (Loo-Choo).  
Distr. Japan (Okinawa-honto, Kume-jima, Miyako-jima, Ishigaki-jima, Iriomote-jima, Iheya-jima).  
*Protaetia (Calopotosia) orientalis* (GORY-PERCHERON, 1833) シロテンハナムグリ  
*Cetonia orientalis* GORY-PERCHERON, 1833, Mon. Ceton., p. 193, tab. 34, fig. 6.  
*Cetonia aerata* ERICHSON, 1834, Nova Acta Halle, XVI, p. 240.  
Distr. Japan, Taiwan, China, Himalaya, N. India, Guamu Is.  
subsp. *submarmorea* (BURMEISTER, 1842) シロテンハナムグリ  
*Cetonia submarmorea* BURMEISTER, 1842, Hand. Entom. III, p. 460.  
Distr. Japan (Hokkaido, Honshu, Tobishima, Awashima, Sado, Izu Is.-Oshima, Hachijyo-Kojima, Hachijyo-jima, Shikoku, Kochi-Okinoshima, Kyushu, Hirato-jima, Danjyo-Gunto, Goto Is., Iki, Tsushima, Hekura-jima, Yakushima, Kuchinoerabu-jima, Amami-Oshima, Okinawa Is., Miyako-jima, Ishigaki-jima), Taiwan.  
subsp. *tokarana* NOMURA, 1964 トカラシロテンハナムグリ  
*Protaetia orientalis tokarana* NOMURA, 1964, Ent. Rev. Japan, XVI, 2, p. 59, tab. 8, fig. 17.  
Distr. Japan (Tokara-Kuchinoshima, Nakanoshima).

Subgenus *Nipponoproteta* MIKESIC, 1966  
*Protaetia* subg. *Nipponoproteta* MIKESIC, 1966, Arbeitsgemein österr. Entom. XVII, 1, p. 66.  
(type-species: *Cetonia lenzi* HAROLD, 1876)

- Protaetia (Nipponoproteta) lenzi* (HAROLD, 1876) レンツアオハナムグリ

*Cetonia lenzi* HAROLD, 1876, Abhand. Nat. Ver. Bremen, V, p. 128.

*Protaetia hondana* ARROW, 1913, Ann. Mag. Nat. Hist. (8):XII, 404.

*Potosia nitidiscutellata* NIIZIMA et KINOSHITA, 1923, Res. Bull. Coll. Exp. Forest Coll. Agr. Hokkaido Imp. Univ., II, 2, pp. 174, 234, taf. V, fig. 2.

Distr. Japan (Honshu, Shikoku, Kyushu, Yakushima).

Subgenus *Pyropotosia* REITTER, 1898

*Pyropotosia* REITTER, 1898, Best.-Taf. XXXVIII, p. 45.

(type-species: *Potosia oschimana* NONFRIED, 1895)

*Protaetia (Pyropotosia) pryeri* (JANSON, 1888) リュキュウツヤハナムグリ

*Cetonia pryeri* JANSON, Mag. Nat. Hist. (6) I : 195, 1888.

Distr. Japan.

subsp. *akuseikiana* NOMURA, 1964 アクセキツヤハナムグリ

*Protaetia pryeri akuseikiana* NOMURA, Tōhō Gakuho (14):68-69, pl. 1, f. 9, 1964 (Akuseki-jima).

Distr. Japan (Akuseki-jima-Tokara Is.).

subsp. *esakii* NAKANE, 1956 エサキツヤハナムグリ

*Protaetia (Pyropotosia) pryeri esakii* NAKANE, Bull. Osaka Munic. Museum. Nat. Hist. (7)7, pl. 1, f. 10, pl. II, f. 6, 1956 (Sata).

Distr. Japan (Sata-Kyushu, Takarajima, Yokoate-jima-Tokara Is.).

subsp. *nitidicosta* YAWATA, 1941 ミヤコツヤハナムグリ

*Protaetia nitidicosta* YAWATA, Kontyu 15(3):122, f. Ka, O. 1941 (Shuri in Okinawa).

Distr. Japan (Miyako-jima, Irabu-jima).

subsp. *oschimana* (NONFRIED, 1895) オオシマツヤハナムグリ

*Potosia oschimana* NONFRIED, 1895, Berliner Entom. Zeitschrift. XL, p. 293.

*Pyropotosia pryeri obenbergeri* BALTHASAR, 1929, Časop. Čsl. Spol. Entom. č. 5-6, p. 114.

*Pyropotosia pryeri* ab. *euchalcea* BALTHASAR, 1929, Acta Soc. ent. Čechoslov. 25, p. 115.

Distr. Japan (Hachijo-jima-Izu Is., Amami-Oshima, Tokunoshima, Kikaigashima).

subsp. *pryeri* (JANSON, 1888) リュウキュウツヤハナムグリ

*Cetonia pryeri* JANSON, 1888, Ann. Mag. Nat. Hist. (6) I, p. 195.

Distr. Japan (Okinoerabu, Iheyajima, Ikinawa Is., Ishigaki-jima).

subsp. *tsutsuii* NAKANE, 1956 トカラツヤハナムグリ

*Protaetia (Pyropotosia) pryeri tsutsuii* NAKANE, Bull. Osaka Munic. Museum. Nat. Hist. (7):6-7, pl. 1, 9, pl. II, f. 5, 1956 (Nakanoshima-Tokara).

Distr. Japan (Kuchinoshima, Nakanoshima-Tokara Is.).

Genus *Gametis* BURMEISTER, 1842 コアオハナムグリ属

*Gametis* (pars A) BURMEISTER, 1842, Handb. Entom. III, p. 356, 358.

*Oxycetonia* ARROW, 1910, Fauna Brit. Ind. Lamell. I, p. 163.

(type-species: *Cetonia versicolor* FABRICIUS, 1775)

*Gametis forticula* (JANSON, 1881) アオヒメハナムグリ

*Glycyphana forticula* JANSON, Cist. Ent. II, 1881, p. 608 (Japan?).

Distr. Japan.

subsp. *forticula* (JANSON, 1881) アオヒメハナムグリ

*Glycyphana forticula* JANSON, Cist. Ent. II, 1881, p. 608 (Japan?).

Distr. Japan (S. Honshu, Shikoku, Kyushu, Tsushima, Hekura-jima, Yakushima, Tokara Is. - Takarajima, Akuseki-jima, Nakanoshima, Amami-Oshima, Tokunoshima, Kikaigashima, Okinawa-to, Kume-jima).

subsp. *ishigakiana* NOMURA, 1959 イシガキヒメハナムグリ

*Oxycetonia jucunda ishigakiana* NOMURA, Tōhō Gakuho (9):52, pl. III, f. 25, 1959 (Ishigaki Is.).

Distr. Japan (Obama-jima, Ishigaki-jima, Taketomi-jima, Iriomote-jima).

subsp. *miyakoana* NOMURA, 1959 ミヤコヒメハナムグリ

*Oxycetonia jucunda miyakoana* NOMURA, Tōhō Gakuho (9):52, 1959 (Miyako Is., Irabu Is.).

Distr. Japan (Miyako-jima, Irabu-jima).

subsp. *yonakuniana* NOMURA, 1959 ヨナクニヒメハナムグリ

*Oxycetonia jucunda yonakuniana* NOMURA, Tōhō Gakuho (9):52, pl. III, f. 26, 1959 (Yonakuni Is.).

Distr. Japan (Yonaguni-jima).

*Gametis jucunda* (FALDERMANN, 1835) コアオハナムグリ

*Cetonia jucunda* FALDERMANN, Ned. Acad. Peterosburg. II, p. 386, pl. IV, f. 4, 5, 1835.

*Gametis argyrosticta* BURMEISTER, Handb. Ent. III, p. 360, 1942.

*Glycyphana albosetosa* MOTSCHULSKY, Et. Ent. X, p. 9, 1861.

*Glycyphana dolens* KRAATZ, Deut. Ent. Zeit. XXIII, p. 236, 1879.

*Gametis feruginosa* REITTER, Verh. Nat. Ver. Brünn. XXXVIII, p. 53, 1898.

Distr. Japan (Hokkaido, Honshu, Tobishima, Awashima, Sado, Izu Is.-Oshima, Toshima, Nii-jima, Shikine-jima, Kozu-jima, Miyake-jima, Hachijyo-jima, Aogashima, Shikoku, Kyushu, Hirato-jima, Goto Is., Iki, Tsushima, Kekura-jima, Tanegashima, Yakushima), Korea, China, Siberia, N. America.

Genus *Callinomes* WESTWOOD, 1874 ホソハナムグリ属

*Callinomes* WESTWOOD, Thes. Ent. Oxon, 1874, p. 26.

(type-species: *Callinomes vollenhovii* WESTWOOD, 1874)

*Callinomes ishikawai* Y. KUROSAWA, 1973 チャイロホソハナムグリ

*Callinomes ishikawai* Y. KUROSAWA, Mem. Nat. Sci. Museum (6):111-112, 1973 (Iriomotejima).

Distr. Japan (Iriomote-jima).

Genus *Coenochilus* SCHAUER, 1841 スジホソハナムグリ属

*Coenochilus* SCHAUER in Germar Zeitschr. Ent. III, 1841, p. 250, 268.

(type-species: *Cetonia maura* FABRICIUS, 1801)

*Coenochilus striatus* WESTWOOD, 1874 スジホソハナムグリ

*Coenochilus striatus* WESTWOOD, Thesaur. Oxon, 1874, p. 46, t. 13, f. 7, 7a-d (Hongkong).

Distr. Japan (Ishigaki Is., Iriomote Is.), Hongkong.